arXiv:quant-ph/9502008v1 8 Feb 1995

Consistent use of paradoxes in deriving constraints on the dynamics of physical systems and of no-go-theorems

K. Svozil
Institut für Theoretische Physik
Technische Universität Wien
Wiedner Hauptstraße 8-10/136
A-1040 Vienna, Austria
e-mail: svozil@tph.tuwien.ac.at

February 1, 2008

Abstract

The classical methods used by recursion theory and formal logic to block paradoxes do not work in quantum information theory. Since quantum information can exist as a coherent superposition of the classical "yes" and "no" states, certain tasks which are not conceivable in the classical setting can be performed in the quantum setting. Classical logical inconsistencies do not arise, since there exist fixed point states of the diagonalization operator. In particular, closed timelike curves need not be eliminated in the quantum setting, since they would not lead to any paradoxical outcome controllability. Quantum information theory can also be subjected to the treatment of inconsistent information in databases and expert systems. It is suggested that any two pieces of contradicting information are stored and processed as coherent superposition. In order to be tractable, this strategy requires quantum computation.

paradox.tex

This letter introduces two novel features of quantum information theory. Physically, it is shown how quantum information allows the consistent implementation of nonlocal correlations. Technically, a diagonalization operator is used to compute consistent fixed point solutions to classical "paradoxical" tasks. The implications for quantum recursion theory [1] and algorithmic information theory [2] as well as for database applications will only be shortly sketched.

Classical information theory (e.g., [3]) is based on the bit as fundamental atom. This classical bit, henceforth called *cbit,* is in one of two classical states.

It is customary to use the symbols "0" and "1" as the names of these states. The corresponding classical bit states are denoted by the symbols 0 and 1.

In quantum information theory (cf. [4, 5, 6, 7, 8, 9, 10, 11]), the most elementary unit of information, henceforth called *qbit*, may be physically represented by a coherent superposition of the two states $|0\rangle$ and $|1\rangle$, which correspond to the symbols 0 and 1, respectively. The quantum bit states

$$|a, b\rangle = a|0\rangle + b|1\rangle \tag{1}$$

form a continuum, with $|a|^2 + |b|^2 = 1$, $a, b \in C$.

In what follows we shall consider the hypothetical transmission of information backward in time. To be more specific, we shall use an EPR-type telegraph which uses entangled particles in a singlet state (i.e., the total angular momentum of the two particles is zero) as drawn in Fig. 1. The apparatus is tuned to convey perfect correlations of the direction of angular momentum labelled by "+" and "−"; i.e., the outcomes are either $++$ or $--$. (Perfect correlations can be achieved by choosing a relative angle of measurement of π.) The (unphysical) assumption necessary for signalling backwards in time is that on one side, say for particles in path 1, the *outcome can be controlled.* This means that it will be assumed possible to produce a particle with, say, direction of angular momentum "+" ("−") in the path 1 at t_A, thereby transmitting a signal "+" ("−") via its perfectly correlated entangled partner in path 2 to a second observer back in time at t_B; thereby, $t_A > t_B > t_S$ but otherwise arbitrary.

An alternative setup for backward in time signalling operates with parameter dependence [12, 13]. There, the (unphysical) assumption is that the measurement outcomes in one path depend on the setting of the measurement angle in the other path.

We shall make use of the EPR-type telegraph to construct a time paradox and argue against parameter dependence [12, 13] and outcome controllability in any form. In a similar manner, the liar paradox [14] was translated by Gödel into arithmetic [15] to argue against a complete description of a formal system within that very system [18]. For instance, the gödelian sentence [19] claiming its own unprovability in a particular system appears undecidable within that very system. In physical terms, undecidability must be translated onto the level of *phenomena.* To put it pointedly: there is no such thing as an inconsistent phenomenon. In a *yes-no* experiment which can have two possible outcomes, only one of these outcomes will actually be measured. There might even be a "hidden parameter (extrinsic [20], exo- [21]) arena," in which this particular outcome could be deterministically accounted for. Yet, for an intrinsic observer who is embedded in the system [22], this level will be permanently inaccessible [23]. As will be shown below, quantum mechanics implements this phenomenological undecidability both by the postulate of randomness of certain outcomes and by the superposition principle. Related arguments have been put forward in [19, 24, 25, 26, 27, 28, 29].



Figure 1: Scheme of backward-in-time signalling by EPR-type telegraph. The postulated controllability of outcomes in 1, mediated via 2, is used to transmit information. The flow of information is indicated by the arrow. "•" stands for the *active* mode; i.e., controllable outcome (preparation). "∘" stands for the *passive* mode; i.e., measurement. The two signs are drawn on top and at bottom to indicate the orientation (relative angle π).

Consider two backward-in-time signalling EPR-type telegraphs of the above type arranged as drawn in Fig. 2. Physically, the flow of information is mediated via the two entangled pairs in paths 1–2 and 3–4. An information in 2 is mirrored by M in 3. By this instrument, some mechanistic agent A (e.g., computer, deterministic observer) which is given the power of outcome control or, alternatively, parameter dependence, can exchange information with itself on closed timelike lines [30, 31, 32, 33]. A shall be confronted with the following paradoxical task. Whenever A registers the information "+" ("−") at time $t_{A'}$, A must stimulate the opposite outcome "−" ("+") at the later time t_A.

Before discussing the paradox, let us consider the two states $|0\rangle \equiv$ "−" and $|1\rangle \equiv$ "+" which are accessible to A. These states can be the basis of a cbit with the identification of the symbols "0" and "1" for $|0\rangle$ and $|1\rangle$, respectively. Quantum mechanically any coherent superposition of them is allowed. A's paradoxical task can be formalized by a unitary evolution operator $\widehat{D}$ as follows

$$\widehat{D}|0\rangle = |1\rangle, \quad \widehat{D}|1\rangle = |0\rangle \quad . \tag{2}$$

In the state basis $\{|0\rangle, |1\rangle\}$, $\widehat{D}$ is just equivalent to the unary logical not-operation and is therefore identical with the not-gate (or the Pauli spin operator τ_1),

$$\widehat{D} = \tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = |1\rangle\langle 0| + |0\rangle\langle 1| \quad . \tag{3}$$

The syntactic structure of the paradox closely resembles Cantor's diagonalization method which has been applied by Gödel, Turing and others for undecidability proofs in a recursion theoretic setup [17, 34, 35, 36]. Therefore, $\widehat{D}$ will be called *diagonalization* operator, despite the fact that its only nonvanishing components are off-diagonal. (Notice that A's task would be perfectly consistent if there were no "bit switch" and if thus $\widehat{D} = \mathrm{diag}(1,1)$.)

The paradoxical feature of the construction reveals itself in the following question: what happens to A? In particular: what does A register and send?

Let us first consider these questions from a classical perspective. Classically, the particles with which A operates can only be in one of two possible states, namely in $|0\rangle$ or in $|1\rangle$, corresponding to the classical bit states. By measuring the particle in beam 4, A gets either the outcome "+" or "−". In both cases, the agent A is lead to a complete contradiction.

For, if A receives "+", corresponding to cbit state 1, A is obliged to send out "−", corresponding to cbit state 0 (A has been assumed to be able to control the outcomes in beam 1). Due to the perfect EPR-correlations, the partner particle in beam 2 is registered as "−" at the mirror at time t_B. By controlling the outcome in beam 3, this mirrored cbit can again be sent backwards in time, where "−" is received by A via a measurement of the particle in beam 4. This, however, contradicts the initial assumption that the outcome in beam 4 is "+".

On the other hand, if A receives "−", corresponding to cbit state 0, A is obliged to send out "+", corresponding to cbit state 1; yet, since at t_B the cbit is



Figure 2: time paradox. Two backward-in-time signalling devices are used here, but only one would be necessary, the other could be subluminal quantum information channel. The important point is the outcome controllability at t_A with regards to the measurement at $t_{A'}$.

just reflected as described above, A should have received "+". Thus classically, agent A is in an inescapable dilemma.

The defense strategy in formal logic and classical recursion theory against such inconsistencies is to avoid the appearance of a paradox by claiming (stronger: requiring) overall consistency, resulting in no-go theorems; i.e., in the postulate of the impossibility of any operational method, procedure or device which would have the potentiality to cause a paradox. (Among the many impossible objects giving rise to paradoxes are such seemingly innocent devices as a "halting algorithm" computing whether or not another arbitrary computable algorithm produces a particular output; or an algorithm identifying another arbitrary algorithm by input-output experiments.)

In the above case, the defense strategy would result in the postulate of the impossibility of any backward-in-time information flow or, more general, of closed timelike lines. Since the only nontrivial feature of the backward-in-time information flow has been outcome controllability or parameter dependence, the diagonalization argument can be used against outcome controllability and parameter dependence, resulting in an intrinsic randomness of the individual outcomes.

Quantum mechanics implements exactly that kind of recursion theoretic argument; yet in a form which is not common in recursion theory. Observe that the paradox is resolved when A is allowed a nonclassical qbit of information. In particular, A's task can consistently be performed if it inputs a qbit corresponding to the *fixed point* state of $\widehat{D}$; i.e.,

$$\widehat{D}|*\rangle = |*\rangle \quad . \tag{4}$$

The fixed point state $|*\rangle$ is just the eigenstate of the diagonalization operator $\widehat{D}$ with eigenvalue 1. Notice that the eigenstates of $\widehat{D}$ are

$$|I\rangle, |II\rangle = \frac{1}{\sqrt{2}}\left[\begin{pmatrix} 1 \\ 0 \end{pmatrix} \pm \begin{pmatrix} 0 \\ 1 \end{pmatrix}\right] = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle) \tag{5}$$

with the eigenvalues $+1$ and -1, respectively. Thus, the nonparadoxical, fixed point qbit in the basis of $|0\rangle$ and $|1\rangle$ is given by

$$|*\rangle = |\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\rangle = |I\rangle \quad . \tag{6}$$

In natural language, this qbit solution corresponds to the statement that it is impossible for the agent to control the outcome, since there is a fifty percent chance for the classical bit states $|0\rangle$ and $|1\rangle$ to be "stimulated" at t_A. The impossibility of outcome control (and parameter independence) is indeed encountered in quantum mechanics [37].

We close the discussion on the consistent use of paradoxes in physics with a few comments. First, it is important to recognize that the above considerations have *no* immediate bearing on quantum complementarity. In the author's opinion, complementarity is a general feature of the intrinsic perception of

computer-generated universes, which is realizable already at a very elementary pre-diagonalization level [38, 39, 36]; i.e., without the requirement of computational universality or its arithmetic equivalent.

As has been pointed out before, the above argument remains valid for any conceivable (local or nonlocal [40, 41]) hidden variable theory. The consistency of the physical phenomenology requires that hidden variables remain inaccessible to an intrinsic observer. Pointedly stated, from an intrinsic, operational point of view, when re-interpreted properly, a paradox marks the appearance of uncertainty and uncontrollability (cf. a statement by Gödel [18]).

A similar treatment of the halting problem [34] for a quantum computer leads to the conclusion that the quantum recursion theoretic "solution" of the halting problem reduces to the tossing of a fair (quantum [42]) coin [43]. Another, less abstract, application for quantum information theory is the handling of inconsistent information in databases. Thereby, two contradicting cbits of information $|a\rangle$ and $|b\rangle$ are resolved by the qbit $|1/\sqrt{2}, 1/\sqrt{2}\rangle = (1/\sqrt{2})(|a\rangle + |b\rangle)$. Throughout the rest of the computation the coherence is maintained [44]. After the processing, the result is obtained by a measurement. The processing of qbits requires an exponential space overhead on classical computers in cbit base [45]. Thus, in order to remain tractable, the corresponding qbits should be implemented on truly quantum universal computers.

Acknowledgements: This and related topics were vigorously discussed in seminars and regular Viennese coffee house sessions with Anton Zeilinger, Johann Sumhammer and others. Günther Krenn took the pain to read, contribute and comment to several versions of the manuscript. An anonymous referee suggested many revisions. Nonetheless, any blame should remain solely with the author.

References

[1] K. Svozil, *Quantum recursion theory*, submitted.

[2] K. Svozil, *Halting probability amplitude of quantum computers*, submitted.

[3] R. W. Hamming, *Coding and Information Theory, Second Edition* (Prentice-Hall, Englewood Cliffs, New Jersey, 1980).

[4] D. Z. Albert, *Phys. Lett.* **94A**, 249 (1983).

[5] D. Deutsch, *Proc. R. Soc. Lond.* **A 400**, 97 (1985).

[6] R. P. Feynman, *Opt. News* **11**, 11 (1985).

[7] A. Peres, *Phys. Rev.* **A32**, 3266 (1985).

[8] P. Benioff, *Annals New York Akademy of Sciences* **480**, 475 (1986).

[9] N. Margolus, *Annals New York Akademy of Sciences* **480**, 487 (1986).

[10] D. Deutsch, *Proc. R. Soc. Lond.* **A 425**, 73 (1989).

[11] D. Deutsch and R. Jozsa, *Proc. R. Soc. Lond.* **A 439**, 553 (1992).

[12] A. Shimony, *Controllable and uncontrollable non-locality*, in *Proc. Int. Symp. Foundations of Quantum Mechanics*, ed. by S. Kamefuchi *et al.* (Physical Society of Japan, Tokyo, 1984), reprinted in A. Shimony, *Search for a Natualistic World View, Volume II* (Cambridge University Press, Cambridge, 1993), p. 130.

[13] A. Shimony, *Events and Processes in the Quantum World*, in *Quantum Concepts in Space and Time*, ed. by R. Penrose and C. I. Isham (Clarendon Press, Oxford, 1986), reprinted in A. Shimony, *Search for a Natualistic World View, Volume II* (Cambridge University Press, Cambridge, 1993), p. 140.

[14] The *Bible* contains a passage, which refers to Epimenides, a Crete living in the capital city of Cnossus: *"One of themselves, a prophet of their own, said, 'Cretans are always liars, evil beasts, lazy gluttons.' "* — St. Paul, Epistle to Titus I (12-13). For more details, see A. R. Anderson, *St. Paul's epistle to Titus*, in *The Paradox of the Liar*, ed. by R. L. Martin (Yale University Press, New Haven, 1970).

[15] K. Gödel, *Monatshefte für Mathematik und Physik* **38**, 173 (1931); English translation in [16] and in Davis, ref. [17].

[16] K. Gödel, *Collected Works, Volume I, Publications 1929-1936*, ed. by S. Feferman, J. W. Dawson, Jr., St. C. Kleene, G. H. Moore, R. M. Solovay, J. van Heijenoort (Oxford University Press, Oxford, 1986).

[17] M. Davis, *The Undecidable* (Raven Press, New York, 1965).

[18] The following citation is taken from K. Gödel's reply to a letter by A. W. Burks, reprinted in J. von Neumann's *Theory of Self-Reproducing Automata*, ed. by A. W. Burks (University of Illinois Press, Urbana, 1966), p. 55; see also S. Feferman, *Philosophia Naturalis* **21**, 546 (1984), p. p. 554:

"... the fact that a complete epistemological description of a language A cannot be given in the same language A, because the concept of truth of sentences of A cannot be defined in A. It is this theorem which is the true reason for the existence of undecidable propositions in the formal systems containing arithmetic."

[19] K. R. Popper, *The British Journal for the Philosophy of Science* **1**, 117, 173 (1950).

[20] K. Svozil, *Europhysics Letters* **2**, 83 (1986); *Il Nuovo Cimento* **96B**, 127 (1986); and [36], chapter 6.

[21] O. E. Rössler, *Endophysics*, in *Real Brains, Artificial Minds*, ed. by J. L. Casti and A. Karlquist (North-Holland, New York, 1987), p. 25; O. E. Rössler, *Endophysics, Die Welt des inneren Beobachters*, ed. by P. Weibel (Merwe Verlag, Berlin, 1992).

[22] T. Toffoli, *The role of the observer in uniform systems*, in *Applied General Systems Research*, ed. by G. Klir (Plenum Press, New York, London, 1978).

[23] The observer is situated, in O. E. Rössler's dictum, in a "Cartesian prison;" cf. H. Atmanspacher and G. Dalenoort, eds., *Inside Versus Outside* (Springer, Berlin, 1994), p. 156.

[24] J. Rothstein, International Journal of Theoretical Physics **21**, 327 (1982).

[25] A. Peres, *Foundation of Physics* **15**, 201 (1985).

[26] St. Wolfram, *Physical Review Letters* **54**, 735 (1985); *Physica* **D10**, 1 (1984); *Rev. Mod. Phys.* **55**, 601 (1983).

[27] Ch. D. Moore, *Phys. Rev. Lett.* **64**, 2354 (1990).

[28] A. C. Elitzur, *Phys. Lett.* **A167**, 335 (1992), as well as S. Popescu and D. Rohrlich, *Foundations of Physics* **24**, 379 (1994) refer to a talk of Professor Y. Aharonov delivered at the Einstein centennial.

[29] A. Posiewnik, *On inconsistency of the language of physics* (University of Gdansk preprint, 1993).

[30] K. Gödel, *Rev. Mod. Phys.* **21**, 447 (1949): K. Gödel, *A remark about the relationship between relativity theory and idealistic philosophy*, in *Albert Einstein, philosopher-scientist*, ed. by P. A. Schilpp (Evanston, Ill., 1949), p. 555.

[31] E. Recami, *Foundations of Physics* **17**, 239 (1987).

[32] J. S. Bell, *Physics* **1**, 195 (1964); J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).

[33] D. Deutsch, *Phys. Rev.* **D44**, 3197 (1991).

[34] H. Rogers, *Theory of Recursive Functions and Effective Computability* (MacGraw-Hill, New York 1967).

[35] P. Odifreddi, *Classical Recursion Theory* (North-Holland, Amsterdam, 1989).

[36] K. Svozil, *Randomness and Undecidability in Physics* (World Scientific, Singapore, 1993).

[37] In an abstract form, the above argument is the analogon to the fixed point or paradoxical combinator in combinatory logic; cf. H. B. Curry and R. Feys, *Combinatory Logic* (North-Holland, Amsterdam, 1958), p. 151, 178 and H. P. Barendregt, *The Lambda Calculus (Revised Edition)* (North Holland, Amsterdam, 1984).

[38] E. F. Moore, *Gedanken-Experiments on Sequential Machines*, in *Automata Studies*, ed. by C. E. Shannon & J. McCarthy (Princeton University Press, Princeton, 1956).

[39] D. Finkelstein and S. R. Finkelstein, *International Journal of Theoretical Physics* **22**, 753 (1983).

[40] I. Pitowsky, *Phys. Rev. Lett.* **48**, 1299 (1982); *Phys. Rev.* **D27**, 2316 (1983); N. D. Mermin, *Phys. Rev. Lett.* **49**, 1214 (1982); A. L. Macdonald, *Ibid.*, 1215 (1982); I. Pitowsky, *Ibid.*, 1216 (1982).

[41] I. Pitowsky, *Quantum Probability − Quantum Logic* (Springer, Berlin, 1989).

[42] K. Svozil, *Phys. Lett.* **140**, 5 (1989).

[43] G. J. Chaitin, *Information, Randomness and Incompleteness, Second edition* (World Scientific, Singapore, 1987, 1990); *Algorithmic Information Theory* (Cambridge University Press, Cambridge, 1987); *Information-Theoretic Incompleteness* (World Scientific, Singapore, 1992).

[44] Note that unlike in fuzzy logic, $|\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\rangle$ is a probability *amplitude* rather than a probability for 0 and 1.

[45] R. P. Feynman, *International Journal of Theoretical Physics* **21**, 467 (1982).